Copyright @ McGraw-Hill Education

nth Roots and Rational Exponents

Explore Inverses of Power Functions

Online Activity Use a calculator to complete the Explore.


INQUIRY What conjectures can you make about $f(x) = x^n$ and $g(x) = \sqrt[n]{x}$ for all odd positive values of n?

Learn *n*th Roots

Finding the square root of a number and squaring a number are inverse operations. To find the square root of α , you must find a number with a square of a. The inverse of raising a number to the nth power is finding the **nth root** of a number. The symbol $\sqrt[n]{}$ indicates an **nth root**.

For any real numbers a and b and any positive integer n, if $a^n = b$, then a is an nth root of b. For example, because $(-2)^6 = 64$, -2 is a sixth root of 64 and 2 is a principal root.

An example of an *n*th root is $\sqrt[4]{36}$, which is read as the nth root of 36. In this example, n is the index and 36 is the radicand, or the expression under the radical symbol.



Some numbers have more than one real nth root. For example, 16 has two square roots, 4 and -4, because 4^2 and $(-4)^2$ both equal 16. When there is more than one real root and n is even, the nonnegative root is called the principal root.

Key Concept • Real nth Roots

Suppose n is an integer greater than 1, a is a real number, and a is an nth root of b.

a	n is even.	n is odd.
<i>a</i> > 0	1 unique positive and 1 unique negative real root: $\pm \sqrt[q]{a}$	1 unique positive and 0 negative real root: $\sqrt[n]{a}$
<i>a</i> < 0	0 real roots	0 positive and 1 negative real root: $\sqrt[p]{a}$
a = 0	1 real root: $\sqrt[n]{0} = 0$	1 real root: $\sqrt[n]{0} = 0$

A radical expression is simplified when the radicand contains no fractions and no radicals appear in the denominator.

Today's Goals

- Simplify expressions involving radicals and rational exponents.
- Simplify expressions in exponential or radical form.

Today's Vocabulary nth root

radicand principal root rational exponent

index

Think About	[t!			
Lorena says she can				
tell that ∛–64 will				
have a real root				

have a real root
without graphing. Do
you agree or disagree?
Explain your reasoning.

Section Company	
	ggerige, grant er sammelien de literate, deline er helmeld in den in den en del bleven mild velocie het versche er en
ı	nganaman ne nemenunun ar andamé e transièté in indifferent all de raphilliain théiligh l'hoanna né a ster, on faoi leis tribia e r
	The second of th
	ggermannings i na inamanana na indhinaka kadinama 1986 i saha sama di ido. Na 1984 andar i na masa mashiri Akk
	$m_{12} = - p_{1}^{2} - m_{2}^{2} m_{1}^{2} m_{2}^{2} m_{3}^{2} m$
	majorani kan kan kan mangangan kalangan kalangan kan majorangan kan kan kan kan kan kan kan kan kan k

Principal Roots

Because $5^2 = 25, 25$ has two square roots, 5 and -5. However, the value of $\sqrt{25}$ is 5 only. To indicate both square roots and not just the principal square root, the expression must be written as $\pm \sqrt{25}$.

Talk About It!

Compare the simplified expressions in the previous example with the ones in this example. Explain why the simplified expressions in this example require absolute value bars when the simplified expressions in the previous example did not.

Simplify.

a. $\pm \sqrt{25x^4}$

$$\pm\sqrt{25x^4} = \pm\sqrt{(\underline{})^2}$$
$$= \pm\underline{}$$

b. $-\sqrt{(v^2+7)^{12}}$

$$-\sqrt{(y^2+7)^{12}} = -\sqrt{[(y^2+7)^6]^2}$$
$$= -(y^2+7)^6$$

c. $\sqrt[3]{343a^{18}b^6}$

$$\sqrt[3]{343a^{18}b^6} = \sqrt[3]{(\underline{})^3}$$

d. $\sqrt{-289c^8d^4}$

There are no real roots of $\sqrt{-289}$. However, there are two imaginary roots, 17i and -17i. Because we are only finding the principal square root, use 17i.

$$\sqrt{-289c^{8}d^{4}} = \sqrt{\underline{} \cdot \sqrt{289c^{8}d^{4}}}$$
$$= \underline{ \cdot \sqrt{289c^{8}d^{4}}}$$

Check

Write the simplified form of each expression.

a.
$$\pm \sqrt{196x^4}$$

a.
$$\pm \sqrt{196x^4}$$
 b. $-\sqrt{196x^4}$ **c.** $\sqrt{-196x^4}$

c.
$$\sqrt{-196x^4}$$

When you find an even root of an even power and the result is an odd power, you must use the absolute value of the result to ensure that the answer is nonnegative.

Example 2 Simplify Using Absolute Value

Simplify.

a. ∜81x⁴

$$\sqrt[4]{81x^4} = \sqrt[4]{(3x)^4} = 3|x|$$

Because x could be negative, you must use the absolute value of x to ensure that the principal square root is nonnegative.

b. $\sqrt[8]{256(y^2-2)^{24}}$

$$\sqrt[8]{256(y^2 - 2)^{24}} = \sqrt[8]{\dots} \cdot \sqrt[8]{(y^2 - 2)^{24}}$$
$$= \underline{\quad | (y^2 - 2)^3 |}$$

Because $(v^2 - 2)^3$ could be negative, you must use the absolute value of $(y^2-2)^3$ to ensure that the principal square root is nonnegative.

Learn Rational Exponents

You can use the properties of exponents to translate expressions from exponential form to radical form or from radical form to exponential form. An expression is in **exponential form** if it is in the form x^n , where n is an exponent. An expression is in radical form if it contains a radical symbol.

For any real number b and a positive integer n, $b^{\frac{1}{n}} = \sqrt[n]{b}$, except where b < 0 and n is even. When b < 0 and n is even, a complex root may exist.

Examples:
$$125\frac{1}{3} = \sqrt[3]{125}$$
 or 5 $(-49)\frac{1}{2} = \sqrt{-49}$ or 7*i*

$$(-49)^{\frac{1}{2}} = \sqrt{-49}$$
 or 7i

The expression $b^{\frac{1}{n}}$ has a **rational exponent**. The rules for exponents also apply to rational exponents.

Key Concept • Rational Exponents

For any nonzero number b and any integers x and y, with y > 1, $b^{\frac{x}{y}} = \sqrt[y]{b^x} = (\sqrt[y]{b})^x$, except when b < 0 and y is even. When b < 0and y is even, a complex root may exist.

Examples:
$$125^{\frac{2}{3}} = (\sqrt[3]{125})^2 = 5^2 \text{ or } 25$$

 $(-49)^{\frac{3}{2}} = (\sqrt{-49})^3 = (7i)^3 \text{ or } -343i$

Key Concept • Simplest Form of Expressions with Rational Exponents An expression with rational exponents is in simplest form when all of the following conditions are met.

- · It has no negative exponents.
- It has no exponents that are not positive integers in the denominator.
- · It is not a complex fraction.
- The index of any remaining radical is the least number possible.

Example 3 Radical and Exponential Forms

Simplify.


a. Write $x^{\frac{4}{3}}$ in radical form.

$$x^{\frac{4}{3}} = \sqrt[3]{x^4}$$

b. Write $\sqrt[5]{x^2}$ in exponential form.

$$\sqrt[5]{x^2} =$$

Go Online You can complete an Extra Example online.



Math History Minute:

Christoff Rudolff (1499-1543) wrote the first German algebra textbook. It is believed that he introduced the radical symbol√ in 1525 in his book Die Coss. Some feel that this symbol was used because it resembled a small r, the first letter in the Latin word radix or root.

Think About It!	
Write two equivalent expressions, one in	
radical form and one in exponential form.	
	endido e territorio esta de esta esta esta esta esta esta esta est
The final standard and we have the standard decision in the standard and the standard and the standard and the standard as the	

to $\frac{1}{4}$?


Think About It!

Why did you set t equal

rational exponents.

Watch Out!

Exponents Recall that when you multiply powers, the exponents are added, and when vou raise a power to a power, the exponents are multiplied.



FINANCIAL LITERACY The expression $c(1 + r)^t$ can be used to estimate the future cost of an item due to inflation, where c represents the current cost of the item, r represents the annual rate of inflation, and t represents the time in years. Write the expression in radical form for the future cost of an item 3 months from now if the annual rate of inflation is 4.7%.

$$c(1+r)^{t} = c(1+0.047)^{\frac{1}{4}}$$
 $r = 0.047, t = \frac{1}{4}$
 $= c(\underline{})^{\frac{1}{4}}$ Add.
 $= c\sqrt[4]{1.047}$ $b^{\frac{1}{n}} = \sqrt[n]{b}$

Example 5 Evaluate Expressions with Rational Exponents

Evaluate $32^{-\frac{2}{5}}$.

$$32^{-\frac{2}{5}} = \frac{1}{32^{\frac{2}{5}}} \qquad b^{-n} = \frac{1}{b^{n}}$$

$$= \frac{1}{(2^{5})^{\frac{2}{5}}} \qquad 32 = 2^{5}$$

$$= \frac{1}{2^{5 \cdot \frac{2}{5}}} \qquad \text{Power of a Power}$$

$$= \text{ or } \frac{1}{4} \qquad \text{Multiply the exponents.}$$

Example 6 Simplify Expressions with Rational Exponents

Simplify each expression.

$$x^{\frac{2}{3}} \cdot x^{\frac{1}{6}} = x^{\frac{2}{3} + \frac{1}{6}}$$
$$= x^{\frac{4}{6} + \frac{1}{6}}$$
$$= x^{\frac{5}{6}}$$

Add powers.

$$\frac{2}{3} = \frac{4}{6}$$

Add the exponents.

b.
$$y^{-\frac{2}{3}}$$

$$y^{-\frac{2}{3}} = \frac{1}{y^{\frac{2}{3}}}$$

$$= \frac{1}{y^{\frac{2}{3}}} \cdot \frac{y^{\frac{1}{3}}}{y^{\frac{1}{3}}}$$

$$= \frac{y^{\frac{1}{3}}}{y^{\frac{3}{3}}} \quad \text{or} \quad \dots$$

$$b^{-n} = \frac{1}{b^n}$$

$$\frac{y^{\frac{1}{3}}}{y^{\frac{1}{3}}} = 1$$

$$y^{\frac{2}{3}} \cdot y^{\frac{1}{3}} = y^{\frac{2}{3}} + \frac{1}{3}$$

 $c. z^{-\frac{1}{3}} \cdot z^{\frac{3}{4}}$ $z^{-\frac{1}{3}} \cdot z^{\frac{3}{4}} = z^{-\frac{1}{3} + \frac{3}{4}}$

Add powers.

$$=$$
 or $z^{\frac{5}{12}}$

$$-\frac{1}{3} = \frac{4}{12}, \frac{3}{4} = \frac{9}{12}$$